

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/26292230>

Terlipressin Continuous Infusion: Please Mind the Solvent!

Article in *Current drug targets* · July 2009

DOI: 10.2174/138945009788488396 · Source: PubMed

CITATIONS

3

READS

3,264

3 authors:



Antonio Rodríguez-Núñez

Complejo Hospitalario Universitario de Santiago and University of Santiago de Com...

454 PUBLICATIONS 5,569 CITATIONS

[SEE PROFILE](#)



Vicente Martínez

Takeda

28 PUBLICATIONS 750 CITATIONS

[SEE PROFILE](#)



Jesús López-Herce

Hospital General Universitario Gregorio Marañón

559 PUBLICATIONS 4,679 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



FAYDS Study [View project](#)



BIOÉTICA UCIP [View project](#)

Terlipressin Continuous Infusion: Please Mind the Solvent!

Antonio Rodriguez-Nunez^{1,*}, Vicente Martínez² and Jesús Lopez-Herce³

¹Pediatric Emergency and Critical Care Division. Hospital Clinico Universitario de Santiago. Santiago de Compostela, Spain; ²Ferring España, SA Medical Department Director. Madrid, Spain and ³Pediatric Intensive Care Unit. Hospital General Universitario Gregorio Marañon. Madrid, Spain

Sir,

Terlipressin (TP) could be an alternative treatment in refractory septic shock [1]. This drug has been used in intravenous bolus, but concerns arise about its side effects, mainly vasoconstriction [2]. Also, bolus may need to be repeated every 4-6 hours and at least one paper indicated a rebound of effects and non-stable clinical situation that might be related with intermittent doses [3].

Although current clinical data are limited and dose ranging studies have not been conducted, continuous TP infusion could be a desirable method of administration in order to titrate effects (both positive and deleterious) [1,4,5].

For continuous TP infusion, the dilution of one vial (1 mg) in 50 ml sterile saline has been recommended. The resultant syringe thus contains 0.02 mg/ml TP and an infusion rate of 1 ml/h represents 0.02 mg/h [1].

This dilution method is simple and similar to the dilution of most of our intensive care drugs but it may not be ideal for TP.

TP is stable only at a narrow pH range (3-4) (manufacturer information). This is not a problem when TP is administered by rapid bolus, but if we make up a dilution with a final pH significantly higher than 4, drug activity may be severely affected.

In order to know what the best dilution for TP is we have measured final pH when different volumes of normal saline vs. 5% glucose are added to prepare a syringe for continuous infusion therapy. The results (Table 1) show that dilutions with normal saline are out of the desired pH range of solution. On the other hand, a simple dilution with 5% glucose maintains the final solution in the optimal pH range.

We consider that this simple but important point must be considered when TP continuous infusion is indicated and prepared. It is possible that published effects of TP continuous infusion are biased by this disturbing factor. Also, this factor should be considered when treatment protocols and clinical trials are designed.

Table 1. Resultant Solution pH after Dilution of One Reconstituted TP Vial in Two Different Solvents

Solution	pH (x + SD)*
Vial per se	3.03 + 0.02
5% glucose (up to 10 ml)	3.24 + 0.01
Normal saline (up to 10 ml)	4.88 + 0.03
5% glucose (up to 50 ml)	3.62 + 0.02
Normal saline (up to 50 ml)	5.50 + 0.01

* The test has been repeated 5 times.

REFERENCES

- [1] Lange M, Ertmer C, Westphal M. Vasopressin vs. terlipressin in the treatment of cardiovascular failure in sepsis. *Intens Care Med* 2008; 34(5): 821-32.
- [2] Rodriguez-Nunez A, Lopez-Herce J, Gil-Anton J, Hernandez A, Rey C, RETSPED Working Group of the Spain Society of Pediatric Intensive Care. Rescue treatment with terlipressin in children with refractory septic shock: a clinical study. *Crit Care* 2006; 10: 1-8.
- [3] Yildizdas D, Yapicioglu H, Celik U, Sertdemir Y, Alhan E. Terlipressin as a rescue therapy for catecholamine-resistant septic shock in children. *Intens Care Med* 2008; 34: 511-17.
- [4] Morelli A, Ertmer C, Lange M, Westphal M. Continuous terlipressin infusion in patients with septic shock: less may be best, and the earlier the better? *Intens Care Med* 2007; 33: 1669-70.
- [5] Umgelter A, Reindl W, Schmid RM, Huber W. Continuous terlipressin infusion in patients with persistent septic shock and cirrhosis of the liver. *Intens Care Med* 2007; 33(9): 1669-70.

*Address correspondence to this author at the Pediatric Emergency and Critical Care Division. Hospital Clinico Universitario de Santiago. Santiago de Compostela, Spain Email: Antonio.Rodriguez.Nunez@sergas.es